

## Elementary Particle Physics: Assignment # 7

Due Thursday April 9th

- 1 The lagrangian for electromagnetic interactions of an electron  $\psi$  (charge -1 and mass  $m$ ) and a scalar  $\phi$  of charge  $e_i$  and mass  $m_s$  with an electric field (photon)  $A$  is

$$\mathcal{L} = \bar{\psi} (i\partial_\mu \gamma^\mu - e\gamma^\mu A_\mu - m) \psi + [(\partial_\mu + iee_i A_\mu)\phi][(\partial^\mu + iee_i A^\mu)\phi]^\dagger - m_s^2 |\phi|^2$$

- With this Lagrangian the amplitude for  $e^-(k, r) + s(p) \rightarrow e^-(k', r') + s(p')$  is

$$M = \frac{e^2 e_i}{q^2} \bar{u}^{r'}(k') (\not{p} + \not{p}') u^r(k)$$

- Obtain the unpolarized squared amplitude and the corresponding differential cross section  $\frac{d\sigma}{dE' d\Omega}$  in the LAB system (where  $p = (m_s, 0)$ ). Neglect the electron mass. As usual  $E'$  and  $\Omega$  are the corresponding energy and solid angle of the outgoing electron.
- With the results above obtain the differential cross section  $\frac{d\sigma}{dE' d\Omega}$  for the DIS  $e^- p \rightarrow e^- X$  in a parton model with partons being scalars.
- Predict the expected scaling and relations between the form factors  $F_1^{ep}$  and  $F_2^{ep}$  in this scalar-parton model