

Unit 13: The Quantum ‘Matrix’

Tzu-Chieh Wei

*C. N. Yang Institute for Theoretical Physics and Department of Physics and Astronomy,
State University of New York at Stony Brook, Stony Brook, NY 11794-3840, USA*

(Dated: October 29, 2024)

In this unit, we discuss quantum simulations and quantum sensing and metrology.

Learning outcomes: You’ll be able to get some glimpses to quantum simulations and quantum sensing/metrology and be able to explain them.

I. INTRODUCTION

Richard Feynman at the annual meeting of APS in 1959 said, “Atoms on a small scale behave like nothing on a large scale, for they satisfy the laws of quantum mechanics. So, as we go down and fiddle around with the atoms down there, we are working with different laws, and we can expect to do different things.” [1]. More than 20 years later in 1981, he gave a lecture on “Simulating physics with computers” and proposed the idea of quantum simulations [2]. He said, “Nature isn’t classical, dammit, and if you want to make a simulation of nature, you’d better make it quantum mechanical, and by golly it’s a wonderful problem, because it doesn’t look so easy.”

He pointed out what we still believe is correct: simulating quantum systems with classical computer requires exponential complexity and proposed to use ‘quantum simulators’ instead for dynamics:

$$|\Psi(t)\rangle = e^{-i \int_0^t dt' H(t')} |\Psi(0)\rangle,$$

where \mathcal{T} denotes the integral is time ordered, i.e., in the discretized version

$$|\Psi(t)\rangle = \lim_{dt \rightarrow 0} e^{H(t-dt)dt} e^{H(t-2dt)dt} \dots e^{H(dt)dt} e^{H(0)dt} |\Psi(0)\rangle.$$

Another topic in this unit is quantum metrology. Heisenberg uncertainty principle [3] is one that is concerning measurement on the position and momentum of a particle, or generally on the two non-commuting observables A and B ,

$$\Delta A \Delta B \geq |\langle \psi | [A, B] | \psi \rangle| / 2.$$

This is where quantum measurement departs from classical measurement. See the link for the historical documents <http://scarc.library.oregonstate.edu/coll/pauling/bond/papers/corr155.1.html> on Heisenberg’s pre-publication proof sheet of his first paper on the uncertainty principle.

II. QUANTUM SIMULATIONS

Classical Church-Turing thesis [1936]. Earlier discussions regarding simulations (of classical systems) were made by Church and Turing, usually referred to as the Church-Turing thesis, which states that, “every function which would naturally be regarded as computable can be computed by the universal Turing machine.” This means that a universal Turing Machine can be used to simulate any other “classical computers”. The thesis also defines the notion of an algorithm that computes a function in terms of such a machine. See Fig. 1 for the illustration of a Turing machine.

Later, David Deutsch was interested in how quantum computers could fit into the statement of Church and Turing, and asked “could the laws of physics be used to derive an even stronger version of the Church–Turing thesis?” What he proposed was that a universal quantum computer (or universal quantum Turing machine) is sufficient to efficiently simulate an arbitrary finite, realizable physical system. This is referred to as the Church-Turing-Deutsch Principle or quantum Church-Turing (Deutsch 1985).

Lloyd’s quantum simulator. In 1996, Seth Lloyd [4] showed that Feynman’s 1982 conjecture is correct that quantum computers can be programmed to simulate any local quantum system (containing few-particle interactions). Lloyd’s idea is to evolve a quantum computer in small time steps to allow efficient simulation of time evolution and that the overall time needed grows only polynomially.

In slightly more details, Lloyd considered a set of Hamiltonians

$$\mathcal{S}_H \equiv \{\tilde{H}_1, \tilde{H}_2, \dots, \tilde{H}_l\},$$




Fig. 3.1 of Nielsen & Chuang

FIG. 1. Illustration of a Turing machine; taken from Fig. 3.6 of the book by Nielsen and Chuang.




FIG. 2. Illustration of local interactions on a lattice.

which can be accessible by turning on and off their action on the time evolution of the quantum system. The unitary operators are thus of the form e^{iAt} , where A is in the algebra generated by \mathcal{S}_H . The mathematical tool is the Suzuki-Trotter decomposition,

$$e^{iHt} = \left(e^{iH_1 t/n} \dots e^{iH_l t/n} \right)^n + \sum_{i>j} [H_i, H_j] t^2 / (2n) + \sum_{k=3}^{\infty} E(k),$$

where $E(k)$ contains higher-order corrections and it is bounded by $\|E(k)\|_{\text{sup}} \leq n \|Ht/n\|_{\text{sup}}^k / k!$ (the supremum is maximum expectation value of the operator over the states of interest). The total error in approximating e^{iHt} is less than $\|n(e^{iHt/n} - 1 - iHt/n)\|_{\text{sup}}$ and can be made as small as desired by taking n sufficiently large. He compared how a quantum simulator works to parking a car, “By going forward and backing up a sufficiently small distance a large enough number of times, it is possible to parallel park in a space only ε longer than the length of the car.”

Let us do some counting. For N_q qubits, the state vector for the system’s wavefunction has 2^{N_q} components and the evolution matrix is of the size $2^{N_q} \times 2^{N_q}$. Thus, to carry out a simulation of a generic quantum system under time evolution by a classical computer, one needs to spend an exponential time complexity to achieve that.

On the other, using one well-controlled quantum system to simulate another or a quantum model, the complexity is only polynomial in time. Let us assume that each H_j in the accessible set acts on at most dimension m , e.g. on two qubits $m = 4$ (see Fig. 2). The number of operations to simulate $e^{iH_j t/n}$ is m^2 at most [5, 6]. Each term is simulated n times and there are l such terms, so the total number of operation is $l n m^2$. For a desired total accumulated error ε , the error in each operation should be less than $\varepsilon / (l n m^2)$. If the system has the typical nearest neighbor or next-nearest neighbor interaction, then $l \sim N_q$. From the expansion (in the Trotter decomposition), the first correction term can give an error $\sim t^2/n$ and we set this to be ε and obtain $n \sim t^2/\varepsilon$ for the simulated time duration t . The duration of each operation is $t/n \sim 1/t$. So the total discrete real time spent is $l(t/n) * n \sim t$. A quantum simulator’s time spent is proportional to the real time duration of the simulated quantum system.

Note in the above we have assumed that the quantum simulator has access to e^{iAt} . If this is not accessible, such

Article

Quantum phase transition from a superfluid to a Mott insulator in a gas of ultracold atoms

Markus Greiner, Olaf Mandel, Tilman Esslinger, Theodor W. Hänsch & Immanuel Bloch

Nature **415**, 39–44 (03 January 2002)

doi:10.1038/415039a

[Download Citation](#)

Received: 26 October 2001

Accepted: 29 November 2001

Published: 03 January 2002




FIG. 3. Illustration of an experiment by Greiner et al. [8].

as in a typical universal quantum computation, where only certain universal gates are available, we need to further decompose e^{iAt} according to the available gate set.

Cirac-Zoller [2012] criteria for quantum simulations. Similar to the DiVincenzo criteria for universal quantum computation, Cirac and Zoller provided a set of desiderata for universal quantum simulations [7].

- **Quantum system** (bosons or fermions with or without spins): it must contain a large number of degrees of freedom. Particles may need to be confined in some region of space. [cf. well-defined two-level systems in DiVincenzo's.]
- **Initialization:** we need to be able to prepare (at least approximately) a known quantum state for the system. [cf. initialized to e.g. zero in DiVincenzo's.]
- **Hamiltonian engineering:** It needs to be possible to engineer a set of interactions with external fields or between different particles, with adjustable values. [cf. universal one- and two-qubit gates in DiVincenzo's.] They may involve a reservoir to simulate open-system dynamics. Among the accessible Hamiltonians there should be some that cannot be efficiently simulated (at present) with classical techniques. [This last one is to ensure certain quantum advantage.]
- **Detection:** We need to be able to perform measurements on the systems. This can be individual (i.e. addressing a few particular sites on the lattice) or collective. Ideally, one should be able to perform single-shot experiments that can be repeated several times; one would be able to determine not only the average of an observable $\langle S \rangle$ but also the average of a function of the observable $\langle f(S) \rangle$. [This corresponds to the readout in DiVincenzo's.]
- **Verification:** Even though there is in no way of efficiently verifying the result if the simulation size is large, one needs to have a way of increasing the confidence in the result. [We do not have a corresponding criterion in DiVincenzo's list.]

How to achieve the last item, verification, is an important question. So far we have very limited tools. We could for example use exactly solvable models in physics to provide a benchmark. We may also use the simulations in forwards and backwards in time to see if the end result is the initial state. Or we could develop more powerful and efficient numerical simulations, such as the Quantum Monte Carlo or tensor network methods.

III. PHYSICAL SYSTEMS AND RECENT EXAMPLES

Superfluid to Mott insulator transition. Arguably the most well-known post-BEC era pioneer paper is the one by M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Quantum phase transition from a superfluid to a Mott insulator in a gas of ultracold atoms, Nature **415**, 39 (2002) [8]; see Fig. 3. The authors trap cold atoms (Rubidium 87) in an optical lattice and vary the trapping depth, which effectively changes the ratio of the tunneling (of a particle from one lattice site to a neighboring one) and the onsite interaction strength t/U . The system is effectively described by the Bose-Hubbard model,

$$H = -t \sum_{\langle i,j \rangle} \hat{b}_i^\dagger \hat{b}_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i,$$

which was proposed and studied theoretically way before the experiment.




FIG. 4. Illustration of an experiment by Jotzu et al. on the topological Haldane model.

Topological Haldane model. In cold atoms, not only bosons can be cooled, fermions can also be used to study low-temperature phases. An experimental paper by G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, *Experimental realization of the topological Haldane model with ultracold fermions*, *Nature* **515**, 237 (2014) [9], cools ^{40}K Potassium atoms trapped in an optical lattice. They realize the so-called topological Haldane model, which is a milestone theoretical work on the Chern insulator and later topological insulators; see Fig. 4. The model is described by the Hamiltonian,

$$H = \sum_{\langle ij \rangle} t_{ij} \hat{c}_i^\dagger \hat{c}_j + \sum_{\langle\langle ij \rangle\rangle} e^{i\Phi_{ij}} t'_{ij} \hat{c}_i^\dagger \hat{c}_j + \Delta_{AB} \sum_{i \in A} \hat{c}_i^\dagger \hat{c}_i.$$

The above two experiments use **ultracold quantum gases**. They have high degree of controllability, novel detection possibilities and extreme physical parameter regimes (compared to solid-state systems); see Bloch, Dalibard & Nascimbène, *Nat. Phys.* 2012 [10]. There are other physical systems, such as trapped ions [see Blatt and Roos, *Nat. Phys.* 2012 [11]], NV diamond centers [12, 13], photonics [14, 15], and superconducting qubits. [16–18].

We list the remaining works mentioned in the lecture:

- “Probing many-body dynamics on a 51-atom quantum simulator,” H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, *Nature* **551**, 579–584 (2017) [19].
- “14-Qubit Entanglement: Creation and Coherence,” T. Monz, P. Schindler, J. T. Barreiro, M. Chwala, D. Nigg, W. A. Coish, M. Harlander, W. Hänsel, M. Hennrich, and R. Blatt, *Phys. Rev. Lett.* **106**, 130506 (2011) [20].
- “Engineered two-dimensional Ising interactions in a trapped-ion quantum simulator with hundreds of spins,” J. W. Britton, B. C. Sawyer, A. C. Keith, C.-C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, *Nature* **484**, 489–492 (2012) [21].
- “Observation of a many-body dynamical phase transition with a 53-qubit quantum simulator,” J. Zhang, G. Pagano, P. W. Hess, A. Kyprianidis, P. Becker, H. Kaplan, A. V. Gorshkov, Z.-X. Gong, and C. Monroe, *Nature* **551**, 601–604 (2017) [22].
- “Observation of a discrete time crystal,” J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, L.-D. Potirniche, A. C. Portter, and C. Monroe, *Nature* **543**, 217–220 (2017) [23].
- “Observation of a discrete time-crystalline order in a disordered dipolar many-body system,” S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, C. von Keyserlingk, N. Y. Yao, E. Demler, and M. D. Lukin, *Nature* **543**, 221–225 (2017) [24].

- “Spectroscopic signatures of localization with interacting photons in superconducting qubits,” P. Poushan, C. Neill, J. Tangpanitanon, V. M. Bastidas, A. Megrant, R. Barends, Y. Chen, Z. Chen, B. Chiaro, et al., J. Martinis, *Science* **358**, 1175-1179 (2017) [25].
- “Quantum supremacy using a programmable superconducting processor,” F. Arute, K. Arya, [...], J. M. Martinis, *Nature* **574**, 505–510 (2019) [26].

A. QuEra’s platform

The company QuEra, created by researchers at Harvard and MIT, has built a 256-atom analog quantum simulator, called Aquila. Their system utilizes Rydberg atoms, which we have explained in an earlier unit. The Hamiltonian is

$$\frac{\mathcal{H}(t)}{\hbar} = \sum_j \frac{\Omega_j(t)}{2} \left(e^{i\phi_j(t)} |0_j\rangle\langle 1_j| + e^{-i\phi_j(t)} |1_j\rangle\langle 0_j| \right) - \sum_j \Delta_j(t) \hat{n}_j + \sum_{j < k} V_{jk} \hat{n}_j \hat{n}_k,$$

where $|0_i\rangle$ is the ground state, $|1_i\rangle$ is some Rydberg state with a large principal quantum number, and $\hat{n}_i = 0|0_i\rangle\langle 0_i| + 1|1_i\rangle\langle 1_i|$ refer to qubits in a position i of space. The interaction term V_{ij} is long-ranged and decays with a power law (6th order) w.r.t. to the distance $|\vec{r}_i - \vec{r}_j|$, and is the reason of the Rydberg blockade mechanism, which induces a positive energy cost if two excitations get too close. In particular,

$$V_{jk} = \frac{C_6}{|\mathbf{x}_j - \mathbf{x}_k|^6},$$

and when two atoms are both in the Rydberg state, the coefficient gets applied where C_6 is the interaction strength, defined as $2\pi \times 862690 \text{ MHz} \cdot \mu\text{m}^6$. Using the Rydberg blockade, it is possible to create a checkerboard pattern of $|0\rangle$ and $|1\rangle$ states (or more precisely, superposition of two possible patterns and perhaps others with much smaller amplitudes).

They allow users to define how many atoms and where they will be located. Then the users design time profiles for the three parameters, Rabi frequency $\Omega(t)$, phase $\phi(t)$ and detuning $\Delta(t)$. At the end, the system measures the excitations (which atom is at which state 0 or 1). At this point, it is still limited and there is no individual addressing, except setting the initial location of atoms. (This in some sense reminds us of D-Wave’s quantum annealers.)

IV. QUANTUM METROLOGY

For two non-commuting observables (Hermitian matrices) A and B , they standard deviations obey

$$\Delta A \Delta B \geq |\langle \psi | [A, B] | \psi \rangle| / 2,$$

where $\Delta A \equiv \sqrt{\langle A^2 \rangle - \langle A \rangle^2}$. The most studied example is the uncertainty between the position x and the momentum p of a particle, $[\hat{x}, \hat{p}] = i\hbar$, which implies that $\Delta x \Delta p \geq \hbar/2$. Heisenberg initially regarded this as a relationship between the precession of a measurement and the disturbance it creates. However, from the derivation, the correct interpretation should be these are intrinsic uncertainties, namely, “if we prepare a large number of quantum systems in identical states, $|\psi\rangle$, and then perform measurements of A on some of those, and of B in others, the statistical uncertainties satisfy above inequality.”

Although Heisenberg’s original idea is intuitive, it is not rigorous. A recent experiment was carried out by Rozema et al. [Phys. Rev. Lett. 109, 100404 (2012)] [27] that uses weak measurement and shows that the original precision-disturbance relationship can be violated. In classical mechanics, measurement of position and momentum or speed can be made separately and arbitrarily precise and there is no fundamental limit. In quantum mechanics, however, the act of measurement changes the system (i.e. some kind of quantum backaction), and thus repeated measurement cannot obtain initial properties, for example, observing the location of a particle ‘localizes’ the particle or collapsing the particle’s wavefunction. That is one reason why the measurement precision and disturbance is not easily quantified simultaneously in an experiment and a weak measurement is needed. But it is possible to formulate some uncertainty relation between measurement precision and disturbance, but one needs to carefully define measurement noise and disturbance and or introduce new inequality, see e.g. Ozawa, *Ann. of Physics*. 2004 [28] and Hofmann, *Phys. Rev. A* 2013 [29].

Deriving the uncertainty principle. Consider two Hermitian operators A and B : $A^\dagger = A$ and $B^\dagger = B$. We rewrite

$$AB = \frac{1}{2}(AB + BA) + \frac{1}{2}(AB - BA) = \frac{1}{2}\{A, B\} + \frac{1}{2}[A, B],$$

evaluate the expectation value with respect to a wavefunction $|\psi\rangle$,

$$\langle\psi|AB|\psi\rangle = \frac{1}{2}\underbrace{\langle\psi|\{A, B\}|\psi\rangle}_{\text{real}} + \frac{1}{2}\underbrace{\langle\psi|[A, B]|\psi\rangle}_{\text{pure imaginary}} =: x + yi.$$

Thus we have

$$\frac{1}{4}|\langle\psi|\{A, B\}|\psi\rangle|^2 + \frac{1}{4}|\langle\psi|[A, B]|\psi\rangle|^2 = |\langle\psi|AB|\psi\rangle|^2 \leq \langle\psi|A^2|\psi\rangle\langle\psi|B^2|\psi\rangle,$$

where we have used the Schwarz inequality $|\vec{a} \cdot \vec{b}|^2 \leq |\vec{a}|^2|\vec{b}|^2$. By replacing A by $A - \langle A \rangle$ and B by $B - \langle B \rangle$, we arrive at

$$\frac{1}{4}|\langle\psi|[A, B]|\psi\rangle|^2 \leq \langle\psi|(A - \langle A \rangle)^2|\psi\rangle\langle\psi|(B - \langle B \rangle)^2|\psi\rangle,$$

which gives the celebrated uncertainty relation.

Standard quantum limit. Let us consider a particle involving in time. Its wavepacket will expand and the uncertainty in its momentum or velocity is

$$\Delta v = \Delta p/m \geq \hbar/(2m\Delta x),$$

and the deviation on the position is

$$\Delta x(t) \sim \Delta x + \frac{\hbar t}{2m\Delta x}.$$

If we minimize this with respect to Δx , we obtain the standard quantum limit,

$$(\Delta x)_{\text{SQL}} = \frac{\hbar t}{2m(\Delta x)_{\text{SQL}}} = \sqrt{\frac{\hbar t}{2m}}.$$

This seems to suggest a limit on how well one can measure the position.

When we consider the phase and number uncertainty, we have

$$[\hat{\phi}, \hat{N}] = i \rightarrow \Delta\phi \geq 1/(2\Delta N) \geq 1/N_{\text{tot}},$$

where in the last inequality we have used the $\Delta N \leq N_{\text{tot}}$. This seems to suggest the limit on the accuracy to measure the phase, which is usually referred to as the Heisenberg limit: the uncertainty in measuring a phase is limited by the total number of photons used.

A natural question is how can we deal with the uncertainty? [See, e.g. Giovannetti, Lloyd & Maccone, Science 06, 1330 (2004) [30].] First, we can monitor only one out of a set of incompatible observables. But this creates large uncertainty in the other noncommuting observable(s). Secondly, we can employ a quantum state in which the uncertainty in the to-be-monitored observable is small (at the cost of a very large uncertainty in complementary observables even before measurement). But there are also errors in addition to the systematic errors. Typical source of errors are environment-induced noise from vacuum fluctuations (the so-called shot noise) and dynamically induced noise in e.g. the position measurement of a free mass (the so-called standard quantum limit). But these do not seem to be a fundamental limit.

Mach-Zehnder interferometer. We have seen the Mach-Zehnder interferometer in Unit 01 in the context of Vaidman's 'interaction-free' measurement. You have also seen that the intensity in the two ports oscillate with the unknown phase, $I_1 = I_0 \cos^2(\phi/2)$, and $I_2 = I_0 \sin^2(\phi/2)$. This can be verified by measuring the intensity, which is equivalent counting the clicks from photo-detectors. If photons are independent then, the uncertainty in the normalized intensity is

$$\Delta(I_1 - I_2)/I_0 = \sin \phi \Delta\phi \propto 1/\sqrt{N}.$$




FIG. 5. Illustration of the Mach-Zehnder interferometer and estimation of a phase ϕ ; see, e.g., https://en.wikipedia.org/wiki/Mach-Zehnder_interferometer.




FIG. 6. Illustration of an experiment by Afek, Ambar & Silberberg, Science 2010 [37], to achieve the Heisenberg limit.

The uncertainty in the phase induced by the sample is limited by the number of photons used (i.e. the shot-noise limit). We will come back to this system later using the quantum Fisher information [31–33].

One natural question is, “Can the shot-noise limit be overcome?” Such a question was studied long ago, see e.g. Caves, PRD 1981 [34]; Barnett, Fabre, Maitre, Eur. Phys. J. D (2003) [35], where it was shown that by using the squeezed vacuum in the original vacuum (no-input) port, one can achieve an uncertainty

$$\Delta\phi \propto 1/N^{3/4}.$$

Later in the development of quantum sensing and metrology, it was shown that this uncertainty can be further improved using entangled states to the Heisenberg limit, i.e., $\Delta\phi \propto 1/N$; see Dowling, arXiv:0904.0163 [36].

The main idea is as follows. If we can create the so-called ‘N00N’ state, which is a two-mode entangled state,

$$|N00N\rangle \equiv (|N_A, 0_B\rangle + |0_A, N_B\rangle)/\sqrt{2}.$$

Then a phase shift in one arm takes the state to

$$(e^{iN\phi}|N_A, 0_B\rangle + |0_A, N_B\rangle)/\sqrt{2}.$$

If one can interfere the two paths, e.g., after beam splitting only measure N -photon detection, then the interference has N times faster oscillation, $|1 + e^{iN\phi}|^2/4$, which can reach the Heisenberg limit; see an experiment in Fig. 6.

Fisher information: classical and quantum. For this part, we refer the readers to Ref. [38]. Let’s begin with the classical part.

Classical part and Cramer-Rao bound. The goal of this discussion is to estimate an unknown parameter θ and to estimate the uncertainty. Given a θ , the measurement that gives some value x is assumed to follow a probability distribution $f(x|\theta)$. From a given reading x_1 , then we estimate the value of θ to be $t(x_1)$. With sufficient statistics, we obtain the exact value

$$\langle t \rangle_\theta = \int dx f(x|\theta) t(x) = \theta.$$




FIG. 7. Illustration of quantum sensing: estimating a phase.

The Fisher information is to quantify how the log likelihood function varies with the unknown parameter,

$$I(\theta) = \int dx f(x|\theta) \left(\frac{\partial}{\partial \theta} \log f(x|\theta) \right)^2 = \left\langle \left(\frac{\partial}{\partial \theta} \log f(x|\theta) \right)^2 \right\rangle_{\theta}.$$

The so-called Cramer-Rao bound on the variance of the estimation t for n independent samples or experiments is given via the Fisher information,

$$\Delta^2 t \geq \frac{1}{nI(\theta)}.$$

Proof.

First, let us consider $n_{\text{sample}} = 1$ and introduce the covariance between two functions $g(x)$ and $h(x)$:

$$\text{Cov}[g, h] \equiv \langle (g - \langle g \rangle)(h - \langle h \rangle) \rangle.$$

Specializing this to the conditions (i) $t(x)$ whose average is $\langle t \rangle = \theta$ and (ii) $\partial L(x|\theta)/\partial \theta$ [with $L(x|\theta) \equiv \log f(x|\theta)$] whose average is 0:

$$\int dx f(x|\theta) \frac{\partial}{\partial \theta} L(x|\theta) = \int dx \frac{\partial}{\partial \theta} f(x|\theta) = \frac{\partial}{\partial \theta} (1) = 0,$$

we have

$$\text{Cov}[t(x), \frac{\partial}{\partial \theta} L(x|\theta)] \equiv \langle (t(x) - \theta)(\frac{\partial}{\partial \theta} L(x|\theta) - 0) \rangle = \langle t(x) \frac{\partial}{\partial \theta} L(x|\theta) \rangle = \int dx t(x) \frac{\partial}{\partial \theta} f(x|\theta) = \frac{\partial}{\partial \theta} \int dx t(x) f(x|\theta) = 1.$$

Then we will use the Cauchy-Schwarz inequality,

$$\Delta^2 g \Delta^2 h = \langle (g - \langle g \rangle)^2 \rangle \langle (h - \langle h \rangle)^2 \rangle \geq |\langle (g - \langle g \rangle)(h - \langle h \rangle) \rangle|^2,$$

noting that

$$\Delta^2 \frac{\partial}{\partial \theta} L = I(\theta),$$

we obtain (for n samples)

$$\Delta^2 t \geq \frac{1}{n_{\text{sample}} I(\theta)}.$$

Quantum part. In this case we use a quantum state ρ that is under some quantum operation Φ_{θ} , as illustrated in Fig. 7, i.e.

$$\rho_{\theta} = \Phi_{\theta}(\rho).$$

We want to estimate the parameter θ and calculate the lower bound on its variance. How do we generalize the classical case? How do we generalize $\partial L(x|\theta)/\partial \theta$ (where $L(x|\theta) \equiv \log f(x|\theta)$)? Intuitively $f(x|\theta) \sim \rho_{\theta}$ (which is defined more precisely being measured w.r.t. an operator Π_x), then

$$\frac{\partial}{\partial \theta} \log f(x|\theta) \sim \frac{\partial \rho_{\theta}}{\partial \theta} / \rho_{\theta} \sim D_{\theta}.$$

Thus, we expect

$$I(\theta) = \left\langle \left(\frac{\partial}{\partial \theta} \log f(x|\theta) \right)^2 \right\rangle_{\theta} \implies \text{Tr}(\rho_{\theta} D_{\theta}^2).$$

It turns out that D_{θ} is defined more precisely via

$$\frac{\partial}{\partial \theta} \rho_{\theta} = \{\rho_{\theta}, D_{\theta}\}/2 =: A,$$

where we also define a quantity A and use the notation of the anticommutator $\{P, Q\} \equiv PQ + QP$.

The Quantum Fisher information is thus

$$I_Q(\rho, \theta) = \text{Tr}(\rho_{\theta} D_{\theta}^2).$$

From this, we expect the corresponding quantum Cramer-Rao bound to be

$$\Delta^2 t \geq \frac{1}{n_{\text{sample}} I_Q(\rho, \theta)}.$$

We will not show the proof, but we note that $f(x|\theta) \equiv \text{Tr}(\rho_{\theta} \Pi_x)$ and the proof should proceed as the classical one.

Unitary process. Let us now consider Φ_{θ} to be a unitary process,

$$\rho_{\theta} = U(\theta) \rho U^{\dagger}(\theta), \quad U(\theta) \equiv e^{-i\theta \hat{G}}.$$

The quantum Fisher information, by definition, is

$$I_Q(\rho, \theta) = \text{Tr}(\rho_{\theta} D_{\theta}^2).$$

From the definition of A , we have

$$A_{ij} = \langle i | \partial \rho_{\theta} / \partial \theta | j \rangle = \langle i | \frac{1}{2} (\rho_{\theta} D_{\theta} + D_{\theta} \rho_{\theta}) | j \rangle,$$

and we can choose specifically $|i\rangle$'s to the eigenbasis, $\rho_{\theta} |j\rangle = \lambda_j |j\rangle$. Thus

$$A_{ij} = \frac{1}{2} (\lambda_i + \lambda_j) D_{ij},$$

which leads to

$$D_{\theta} = 2 \sum_{i,j} \frac{\langle i | \partial \rho_{\theta} / \partial \theta | j \rangle}{\lambda_i + \lambda_j} |i\rangle \langle j|.$$

In our case study where Φ_{θ} is unitary, the eigenspectrum of ρ_{θ} is the same as that of ρ , and we have that the Quantum Fisher information is independent of θ , (noting $\partial \rho_{\theta} / \partial \theta = -i[\hat{G}, \rho_{\theta}]$)

$$I_Q(\rho, \theta, \hat{G}) = 2 \sum_{i,j} \frac{(\lambda_i - \lambda_j)^2}{\lambda_i + \lambda_j} |\langle i_0 | G | j_0 \rangle|^2, \quad \rho = \sum_i \lambda_i |i_0\rangle \langle i_0|,$$


where the subscript 0 is used to denote the original eigenbasis. The detailed derivation is left as an exercise, but we have the following,

$$I_Q(|\psi\rangle, \theta, \hat{G}) = 4(\langle \psi | \hat{G}^2 | \psi \rangle - \langle \psi | \hat{G} | \psi \rangle^2) = 4(\Delta G)^2,$$

$$\Delta^2 t \geq \frac{1}{n_{\text{sample}} I_Q(\rho, \theta)}.$$

Example. Coherent state: We have seen the coherent state in an earlier unit, which is a good description of lasers.

$$\rho = |\alpha\rangle \langle \alpha|, \quad \hat{a} |\alpha\rangle = \alpha |\alpha\rangle$$



[e.g. see Tan & Jeong, arXiv: 1909.00942]

FIG. 8. Illustration of an interferometer and estimation of phases; Tan and Jung, arXiv:1909.00942 [38].

We want to see how good it is for the estimation of a unknown phase. Plugging it into the Fisher information, we find that $I_Q = |\alpha|^2$ is proportional to the average number of photons.

Fock states: these are fixed photon-number states,

$$\rho = |n\rangle\langle n|, \quad \hat{a}|n\rangle = \sqrt{n}|n-1\rangle.$$

It is easy to see that

$$I_Q = 0,$$

and hence these states are not useful for estimating the phase, as it will gives very large uncertainty, $\Delta t \sim 1/\sqrt{I_Q}$.

Superposition: What if we consider a superposition of two number states, say, the vaccum and $|n\rangle$?

$$\rho = |\psi\rangle\langle\psi|, \quad |\psi\rangle = (|0\rangle + |n\rangle)/\sqrt{2}$$

It is straightforward to show that

$$I_Q = n^2,$$

which achieves the Heisenberg limit, as $\Delta t \sim 1/n$.

When the input is pure and there are multiple parameters. We have seen that for a single parameter case,

$$|\psi_\theta\rangle = U(\theta)|\psi\rangle, \quad |\partial_\theta\psi_\theta\rangle \equiv \frac{\partial}{\partial\theta}|\psi_\theta\rangle,$$

the quantum Fisher information for a pure state has a much simpler expression:

$$I_Q(|\psi\rangle, \theta) = 4(\langle\partial_\theta\psi_\theta|\partial_\theta\psi_\theta\rangle - |\langle\psi_\theta|\partial_\theta\psi_\theta\rangle|^2).$$

We can generalize this to multiple parameters and define the following quantum Fisher ‘matrix’:

$$(I_Q)_{i,j} = 4\text{Re}(\langle\partial_{\theta_i}\psi_\theta|\partial_{\theta_j}\psi_\theta\rangle - \langle\psi_\theta|\partial_{\theta_i}\psi_\theta\rangle\langle\partial_{\theta_j}\psi_\theta|\psi_\theta\rangle).$$

We note that the quantum Fisher ‘matrix’ (or ‘metric’) was recently used in the gradient decent approach for the VQE algorithm. See e.g. “Simultaneous Perturbation Stochastic Approximation of the Quantum Fisher Information,” by Julien Gacon, Christa Zoufal, Giuseppe Carleo, Stefan Woerner, Quantum 5, 567 (2021) at <https://quantum-journal.org/papers/q-2021-10-20-567/> [39] and “Fisher Information in Noisy Intermediate-Scale Quantum Applications,” by Johannes Jakob Meyer, published in Quantum 5, 539 (2021), <https://arxiv.org/pdf/2103.15191.pdf> [40], with a perspective article by K. Bharti at <https://quantum-journal.org/views/qv-2021-10-06-61/> [41].

Example. See Fig. 8 for the illustration.

$$\hat{G} = (\hat{a}^\dagger\hat{a} - \hat{b}^\dagger\hat{b})/2$$

$$I_Q(|\psi\rangle, \theta, \hat{G}) = 4(\langle\psi|\hat{G}^2|\psi\rangle - \langle\psi|\hat{G}|\psi\rangle^2) = 4(\Delta G)^2.$$

There are two modes, and \hat{G} introduces a phase difference between the two modes. We will send separately coherent states into these,

$$|\alpha\rangle \otimes |\beta\rangle, \quad \hat{a}|\alpha\rangle = \alpha|\alpha\rangle, \quad \hat{b}|\beta\rangle = \beta|\beta\rangle.$$

We find that the use of coherent states (in two modes) leads to the shot-noise limit,

$$I_Q = |\alpha|^2 + |\beta|^2.$$

However, if we send in a N00N state:

$$|\text{N00N}\rangle \equiv (|N_A, 0_B\rangle + |0_A, N_B\rangle)/\sqrt{2},$$

we find that

$$I_Q = N^2,$$

showing that we can achieve the Heisenberg limit.

V. POTENTIAL APPLICATIONS

Waveform of a weak classical force. The fundamental limit of detecting random forces needs to account for the quantum nature of measurement; see a recent paper for further discussions [42].

Gravitational waves. In recent years, we have witnessed advancement in LIGO's experiments that have detected many events generating gravitational waves. These are a feat of metrology (and physics and engineering) on their own. LIGO was using coherent light (lasers) in the Michelson-Morley interferometer setup and has recently moved to the use of squeezed states (which we probably have not actually defined in this course) to enhance its detection sensitivity so as to detect fainter signals.

We learned previously that a coherent state $|\alpha\rangle$ is an eigenstate of an annihilation operator \hat{a} : $\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$, and the coherent state is an infinite superposition of number states, i.e.,

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} |n\rangle.$$

Its standard deviation of any two conjugated quadratures analogous to x and p is equal to each other and saturates the uncertainty relation $\Delta x \Delta p = \hbar/2$. A one-mode squeezed state is

$$|\xi\rangle = e^{\xi^* \hat{a}^2/2 - \xi(\hat{a}^\dagger)^2/2} |0\rangle.$$

This squeezed vacuum is composed of an infinite superposition of even number states and has one quadrature with $\Delta q < \sqrt{\hbar/2}$ (depending on the phase of the complex number ξ), therefore the name squeezed. Given the smaller Δq , then it is possible to achieve better measurement than the coherent state for certain observables. There is a news about "LIGO surpasses the quantum limit" <https://www.ligo.caltech.edu/news/ligo20231023>, which I assume is the standard quantum limit we discussed earlier. One of the earliest papers is Ref. [43].

Dark matters? Another great interest in applying quantum metrology is the detection of the dark matter. Surprising, this unknown matter consists of about 26.8% of the whole universe (in terms of energy) and, in contrast, the ordinary matter (that we and our world are made of) is only about 5%. Unfortunately, dark matter does not interact with the ordinary matter, except through gravity (which is how it was discovered). It is thus not known how to detect them in labs. There are many theoretical proposals of what could constitute dark matter, such as weakly interacting massive particles (WIMPs), sterile neutrinos (a forth one beyond the known three associated with electrons, muons and taus) and axions. Different types of dark matter likely require different technology of detection. Could quantum information and technology help? Researchers have begun investigating whether quantum states of light can help; for example, Backes et al. found that squeezed states of light may enhance search for dark matter axions [44]. Another group of scientists (Dixit et al.) found that using superconducting qubits, it is possible to enhance the low mass bosonic dark matter (such as the axion or hidden photon) by a factor of 1,300 [45]. It is expected more works using quantum metrology for dark matter search will appear and one hopes that dark matter could be found and studied in the lab someday.

VI. CONCLUDING REMARKS

In this unit, we have discussed quantum simulations and quantum sensing and metrology.

It is a good time to check whether you have achieved the following Learning Outcomes:

After this Unit, You'll be able to get some glimpses to quantum simulations and sensing and metrology and to explain them.

Suggested reading: Richard Feynman's paper "Simulating Physics with Computers", Int. J. Theor. Physics, vol 21, no.6/7, p467-488 (1982) [2].

-
- [1] R. P. Feynman, "there's plenty of room at the bottom". engineering and science (california institute of technology) (1960).
 - [2] R. P. Feynman, Simulating physics with computers 1981, International Journal of Theoretical Physics **21**, 467 (1982).
 - [3] W. Heisenberg, Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik, Zeitschrift für Physik **43**, 172 (1927).
 - [4] S. Lloyd, Universal quantum simulators, Science **273**, 1073 (1996).
 - [5] S. Lloyd, Almost any quantum logic gate is universal, Physical review letters **75**, 346 (1995).
 - [6] D. E. Deutsch, A. Barenco, and A. Ekert, Universality in quantum computation, Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences **449**, 669 (1995).
 - [7] J. I. Cirac and P. Zoller, Goals and opportunities in quantum simulation, Nature physics **8**, 264 (2012).
 - [8] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Quantum phase transition from a superfluid to a mott insulator in a gas of ultracold atoms, nature **415**, 39 (2002).
 - [9] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, Experimental realization of the topological haldane model with ultracold fermions, Nature **515**, 237 (2014).
 - [10] I. Bloch, J. Dalibard, and S. Nascimbene, Quantum simulations with ultracold quantum gases, Nature Physics **8**, 267 (2012).
 - [11] R. Blatt and C. F. Roos, Quantum simulations with trapped ions, Nature Physics **8**, 277 (2012).
 - [12] L. Childress and R. Hanson, Diamond nv centers for quantum computing and quantum networks, MRS bulletin **38**, 134 (2013).
 - [13] V. Dobrovitski, G. Fuchs, A. Falk, C. Santori, and D. Awschalom, Quantum control over single spins in diamond, Annu. Rev. Condens. Matter Phys. **4**, 23 (2013).
 - [14] P. Kok, W. J. Munro, K. Nemoto, T. C. Ralph, J. P. Dowling, and G. J. Milburn, Linear optical quantum computing with photonic qubits, Rev. Mod. Phys. **79**, 135 (2007).
 - [15] S. Slussarenko and G. J. Pryde, Photonic quantum information processing: A concise review, Applied Physics Reviews **6**, 041303 (2019).
 - [16] K. Osbourne and J. Martinis, Superconducting qubits and the physics of josephson junctions, in *Les Houches conference proceedings* (2003).
 - [17] J. M. Gambetta, J. M. Chow, and M. Steffen, Building logical qubits in a superconducting quantum computing system, npj quantum information **3**, 1 (2017).
 - [18] P. Krantz, M. Kjaergaard, F. Yan, T. P. Orlando, S. Gustavsson, and W. D. Oliver, A quantum engineer's guide to superconducting qubits, Applied Physics Reviews **6**, 021318 (2019).
 - [19] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, *et al.*, Probing many-body dynamics on a 51-atom quantum simulator, Nature **551**, 579 (2017).
 - [20] T. Monz, P. Schindler, J. T. Barreiro, M. Chwalla, D. Nigg, W. A. Coish, M. Harlander, W. Hänsel, M. Hennrich, and R. Blatt, 14-qubit entanglement: Creation and coherence, Phys. Rev. Lett. **106**, 130506 (2011).
 - [21] J. W. Britton, B. C. Sawyer, A. C. Keith, C.-C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, Engineered two-dimensional ising interactions in a trapped-ion quantum simulator with hundreds of spins, Nature **484**, 489 (2012).
 - [22] J. Zhang, G. Pagano, P. W. Hess, A. Kyprianidis, P. Becker, H. Kaplan, A. V. Gorshkov, Z.-X. Gong, and C. Monroe, Observation of a many-body dynamical phase transition with a 53-qubit quantum simulator, Nature **551**, 601 (2017).
 - [23] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I.-D. Potirniche, A. C. Potter, A. Vishwanath, *et al.*, Observation of a discrete time crystal, Nature **543**, 217 (2017).
 - [24] S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, *et al.*, Observation of discrete time-crystalline order in a disordered dipolar many-body system, Nature **543**, 221 (2017).
 - [25] P. Roushan, C. Neill, J. Tangpanitanon, V. M. Bastidas, A. Megrant, R. Barends, Y. Chen, Z. Chen, B. Chiaro, A. Dunsworth, *et al.*, Spectroscopic signatures of localization with interacting photons in superconducting qubits, Science **358**, 1175 (2017).
 - [26] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. Brandao, D. A. Buell, *et al.*, Quantum supremacy using a programmable superconducting processor, Nature **574**, 505 (2019).

- [27] L. A. Rozema, A. Darabi, D. H. Mahler, A. Hayat, Y. Soudagar, and A. M. Steinberg, Violation of heisenberg's measurement-disturbance relationship by weak measurements, *Phys. Rev. Lett.* **109**, 100404 (2012).
- [28] M. Ozawa, Uncertainty relations for noise and disturbance in generalized quantum measurements, *Annals of Physics* **311**, 350 (2004).
- [29] H. F. Hofmann, Uncertainty characteristics of generalized quantum measurements, *Phys. Rev. A* **67**, 022106 (2003).
- [30] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum-enhanced measurements: beating the standard quantum limit, *Science* **306**, 1330 (2004).
- [31] C. W. Helstrom, Quantum detection and estimation theory, *Journal of Statistical Physics* **1**, 231 (1969).
- [32] V. Giovannetti, S. Lloyd, and L. Maccone, Advances in quantum metrology, *Nature photonics* **5**, 222 (2011).
- [33] G. Tóth and I. Apellaniz, Quantum metrology from a quantum information science perspective, *Journal of Physics A: Mathematical and Theoretical* **47**, 424006 (2014).
- [34] C. M. Caves, Quantum-mechanical noise in an interferometer, *Phys. Rev. D* **23**, 1693 (1981).
- [35] S. M. Barnett, C. Fabre, and A. Maitre, Ultimate quantum limits for resolution of beam displacements, *The European Physical Journal D-Atomic, Molecular, Optical and Plasma Physics* **22**, 513 (2003).
- [36] J. P. Dowling, Quantum optical metrology – the lowdown on high-n00n states, *Contemporary Physics* **49**, 125 (2008), <https://doi.org/10.1080/00107510802091298>.
- [37] I. Afek, O. Ambar, and Y. Silberberg, High-noon states by mixing quantum and classical light, *Science* **328**, 879 (2010).
- [38] K. C. Tan and H. Jeong, Nonclassical light and metrological power: An introductory review, *AVS Quantum Science* **1**, 014701 (2019).
- [39] J. Gacon, C. Zoufal, G. Carleo, and S. Woerner, Simultaneous perturbation stochastic approximation of the quantum fisher information, *Quantum* **5**, 567 (2021).
- [40] J. J. Meyer, Fisher information in noisy intermediate-scale quantum applications, *Quantum* **5**, 539 (2021).
- [41] K. Bharti, Fisher information: A crucial tool for nisc research, *Quantum Views* **5**, 61 (2021).
- [42] J. W. Gardner, T. Gefen, S. A. Haine, J. J. Hope, J. Preskill, Y. Chen, and L. McCuller, Stochastic waveform estimation at the fundamental quantum limit, *arXiv preprint arXiv:2404.13867* (2024).
- [43] J. Aasi, J. Abadie, B. Abbott, R. Abbott, T. Abbott, M. Abernathy, C. Adams, T. Adams, P. Addesso, R. Adhikari, *et al.*, Enhanced sensitivity of the ligo gravitational wave detector by using squeezed states of light, *Nature Photonics* **7**, 613 (2013).
- [44] K. M. Backes, D. A. Palken, S. A. Kenany, B. M. Brubaker, S. Cahn, A. Droster, G. C. Hilton, S. Ghosh, H. Jackson, S. K. Lamoreaux, *et al.*, A quantum enhanced search for dark matter axions, *Nature* **590**, 238 (2021).
- [45] A. V. Dixit, S. Chakram, K. He, A. Agrawal, R. K. Naik, D. I. Schuster, and A. Chou, Searching for dark matter with a superconducting qubit, *Physical review letters* **126**, 141302 (2021).